
NAVAL POSTGRADUATE SCHOOL MONTEREY CA F/6 8/3 ORGANIZATION OF AN OCEANDGRAPHIC DATA BANK FOR THE PERUVIAN NAV--ETC(U) SEP 81 H S SOLDI AD-A114 548 UNCLASSIFIED Ins # 4 4546 (3) END DATE 6-82 DTIC



NAVAL POSTGRADUATE SCHOOL Monterey, California



THESIS

ORGANIZATION OF AN OCEANOGRAPHIC DATA BANK FOR THE PERUVIAN NAVY

by

Hector Soldi Soldi

September 1981

Thesis Advisor:

R. Garwood

Co-Advisor:

D. McLain

Approved for public release, distribution unlimited

S MAY 1 9 1982

82 05 17 030

LILLE FILE COP

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM	
AD-A114548	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitio)	S. TYPE OF REPORT & PERIOD COVERED	
Organization of an Oceanographic Data Bank for	Master's Thesis September 1981	
the Peruvian Navy	E. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(e)	8. CONTRACT OR GRANT NUMBER(s)	
Hector Soldi Soldi		
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
Naval Postgraduate School Monterey, California 93940		
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE	
Naval Postgraduate School	September 1981	
Monterey, California 93940	80	
14. MONITORING AGENCY NAME & ADDRESS(If different from Controlling Office)	15. SECURITY CLASS, (of this report)	
	Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING	
IG. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release, distribution unlimited	i	
17. DISTRIBUTION STATEMENT (of the electroct entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) OCEANOGRAPHIC data, Peru Current, "El Nino", data exchange, data bank, Peruvian Navy, oceanographic data acquisition, Peru		
Oceanographic data have been acquired along the coast of Peru for several decades. This information is important due to the effect of the cold northward Peru Current and Undercurrent and the intermittent warm "El Nino" countercurrent along the coast of Peru. These two phenomena greatly affect the fishing grounds along the coast as well as the characteristics of sound propagation in the sea due to changes in the vertical thermal structure of the water column. The creation of an oceanographic data bank for the Peruvian Navy is proposed for the archival from various sources of all available historical data for the		

SECURITY CLASSIFICATION OF THIS PAGETHEN Role Batome

waters near Peru. The data bank will be updated by a long term program of oceanographic data acquisition and exchange with local and foreign institutions. These data will be stored in an easily accessible format that can be useful for the Navy in creating operational products for the fleet and in monitoring and eventually predicting the ocean variability along the coast of Peru.

Some of the applications of the data bank for the Navy, fisheries and other potential users are presented. $_{\rm A}$

DTIC COPY INSPECTED 2

Accession For 🤏
PTIS WASI
INIT TAB
U.Managapaced [1]
Justification
} B♥
Piters butlen/
Availab lity Codes
tiens and ten
Mat 1 English
101
Manager designation to the contraction of the contr

Approved for public release, distribution unlimited

Organization of an Oceanographic Data Bank for the Peruvian Navy

bу

Hector Soldi Soldi Lieutenant, Peruvian Navy B.A., Peruvian Naval Academy, 1973

Submitted in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE IN OCEANOGRAPHY

from the

NAVAL POSTGRADUATE SCHOOL September 1981

Author:	Commence of the Commence of th
Assumed him	Roland W. Samood 12
Approved by: _	Thesis Advisor
	Douglas R. McLain
-	Co-Advisor
-	Chairman, Department of Oceanography
-	Dean of Science and Engineering

ABSTRACT

Oceanographic data have been acquired along the coast of Peru for several decades. This information is important due to the effect of the cold northward Peru Current and Undercurrent and the intermittent warm "El Nino" countercurrent along the coast of Peru. These two phenomena greatly affect the fishing grounds along the coast as well as the characteristics of sound propagation in the sea due to changes in the vertical thermal structure of the water column. The creation of an oceanographic data bank for the Peruvian Navy is proposed for the archival from various sources of all available historical data for the waters near Peru. The data bank will be updated by a long term program of oceanographic data acquisition and exchange with local and foreign institutions. These data will be stored in an easily accessible format that can be useful for the Navy in creating operational products for the fleet and in monitoring and eventually predicting the ocean variability along the coast of Peru.

Some of the applications of the data bank for the Navy, fisheries and other potential users are presented.

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude to my advisors, Dr. Roland W. Garwood, and Dr. Douglas McLain, for their assistance, and to the Fleet Numerical Oceanography Center at Monterey for supplying the data for this work.

TABLE OF CONTENTS

ABS	TRACT	「	3
ACK	NOWL	EDGEMENTS	4
I.	INT	RODUCTION	9
II.	OCEA	ANOGRAPHIC DATA FROM PERUVIAN WATERS	12
	Α.	HISTORICAL REMARKS	12
	В.	OCEANOGRAPHIC EXPEDITIONS	14
		1. EASTROPAC Program	16
		2. El Nino Watch Expedition	17
		3. CUEA Program	17
III.	THE	IMPORTANCE OF AN OCEANOGRAPHIC DATA BANK IN PERU	20
	Α.	THE FISHERIES	20
	В.	THE NAVY	23
	c.	THE CREATION OF THE BANK	24
IV.	ORG	ANIZATION OF THE BANK	28
	Α.	THE IMPORTANCE OF DATA FORMATS	28
	В.	HISTORICAL DATA FORMAT (MOODS)	31
	c.	DESCRIPTION OF THE MOODS FORMAT	32
		1. Day-Time Group	32
		2. Position	32
		3. Cruise Number and Ship Code	32
		4. Number of Levels	33
		5. Number of Standard Levels	33
		6. Report Number	33

		7. Profile Data	33
		8. Source Code	34
	D.	EXTRACT FILE FOR THE PERUVIAN AREA	34
	Ε.	SELECTION OF THE DATA	37
٧.	APP	LICATIONS	38
	Α.	INTRODUCTION	38
	В.	INVENTORY OF THE DATA	39
	c.	NAVAL APPLICATIONS	47
	D.	OTHER APPLICATIONS	49
		1. Topography of Isothermal Surfaces	49
		2. Sea Level Data	54
VI.	CON	CLUSIONS	55
APP	ENDI	X A - PRINT FORMAT OF HYDRO-CAST AND BT PROFILES	57
APP	ENDI	X B - LISTING OF OBSERVATIONS	58
LIS	T OF	REFERENCES	74
TNT	TTAI	DISTRIBUTION LIST	76

LIST OF TABLES

١.	Data Collection Programs	1950's-1970's	15
2.	Flow of Information into	the Bank	27

LIST OF FIGURES

1.	Area Selected for Data Extraction from MOODS Monthly Tapes	35
2.	Distribution of BT Historical Data for 1960	41
3.	Distribution of Hydrocast Historical Data for 1960	42
4.	Distribution of Hydrocast and BI Data for 1960	43
5.	BT Historical Data for the Month of June, All Years	44
6.	Hydrocast Historical BT Data for the Month of June, All Years	45
7.	Hydrocast and Historical BT Data for the Month of June, All Years	46
8.	Map with Grid of One Degree Quadrangles	48
9.	Sound Speed Profiles for Period April-June Quadrange #60	50
10.	Trans. Loss Curves for Representative Profile at 6 Frequencies	51
11.	Map of Depth(m) of 15°C Isotherm for May 1971	53

I. INTRODUCTION

Oceanographic data have been acquired along the coast of Peru for many decades. The special characteristics of the Peruvian coastal waters, especially the strong upwelling of cold, nutrient rich waters along the coast and the sudden and sometimes catastrophic "El Nino" phenomenon, have attracted the interest of ocean scientists for many years.

The economic, social and environmental impact of these two features on Peruvian society, and on the whole world, has motivated the organization of several large oceanographic expeditions to the coast of Peru. Due to the extent and the diversity of information acquired, the expeditions of international cooperative programs that have taken place during the last fifteen years are most important.

Unfortunately, much of the data that has been collected at high cost and effort for so many years is not centralized and is, therefore, not readily available to potential users in Peru or elsewhere. This is particularly true for physical oceanographic data, since much biological and chemical data have been assembled by the fisheries institutes.

The Peruvian Navy requires an easily accessible oceanographic data bank to be able to provide information to its ships and submarines about ocean conditions under which they operate, and to make ocean forecasts for naval operation areas. However, the task of assembling a data bank is difficult because many of the data are not available in a computer accessible form and because no single institution or agency in Peru has

archaved all of the available historical data. In addition, the advantages of having an oceanographic data bank in Peru will benefit the fisheries, oceanographers, and scientific research in general, as well as the Navy.

In 1973, the Naval Hydrographic Office (Direction de Hidrografia y Navegacion de la Marina--DHNM) was assigned the task of implementing a "System of Oceanographic Information" (Sistema de Informacion Oceanographico, or SIO). This system was envisioned as consisting of an oceanographic data base that would provide services to several users. There would need to be a long term program of oceanographic data acquisition to update the data bank on a continuing basis. The system would also have a mechanism for exchanging data with other local and foreign institutions.

This thesis is a pilot study for the organization of a data bank for the Peruvian Navy that will eventually be part of the SIO. The scope of this work has been focused on the assembly of all available historical oceanographic data for areas in the vicinity of Peru. The historical data was mainly extracted from the Master Oceanographic Observation Data Set (MOODS) files of the Fleet Numerical Oceanographic Center (FNOC) in Monterey, California. After the data were selected, a simple system was designed to identify and store them in an orderly manner.

This thesis has been divided into six chapters. The first chapter is an introduction. The second is an historical review of the data acquisition programs that have taken place off the coast of Peru with a brief description of the most important. The third chapter reviews the phenomena of upwelling and "El Nino" and their impact on the guano and

fisheries industry in Peru, pointing out the importance of oceanographic data for study, monitoring and eventually predicting such features. It also presents justification for having a long range program of data acquisition to update the data bank. The fourth chapter introduces the MOODS system which was used in the selection and processing of the data, commenting on the advantages and disadvantages of implementing a similar system for the SIO. The fifth chapter presents some examples of practical applications of the data and discusses future possible uses of the bank. The last chapter presents the conclusions.

II. OCEANOGRAPHIC DATA FROM PERUVIAN WATERS

A. HISTORICAL REMARKS

Ever since 1811 when A. Von Humboldt described [Humboldt, 1826] what is now called the Humboldt or Peru current, many scientists and investigators have studied the coastal waters of Peru. It is difficult to identify when the peculiar characteristics of the cool waters of the Peruvian coast first drew the attention of the people living or traveling along this narrow strip of land. However, there are several Spanish documents dated as early as 1540, issued to the King of Spain, that report [Murphy, 1937] on the geographical and climatic characteristics of the newly conquered land. The cool ocean waters, the almost total lack of rain, and the abundance of marine life were the most striking features for the newcomers, who reported how they cooled their drinks by hanging flasks over the side of their ships.

One of the most notorious characteristics that concerned the Spanish, however, (because of its implications for navigation) was the origin and path of the sometimes strong northward current that affected the route of the ships sailing between Panama and Callao. An excerpt [Zarate, 1555], from the "Account of the Country" written by Treasurer-General Zarate to the Spanish King in 1543, indicates their preoccupation with this fact:

"This constant wind and current render the navigation exceedingly difficult from Panama to Peru for the greater part of the year; so that vessels are obliged always to tack to windward against wind and current."

Since the time of the Spanish conquest, many other travelers, scientists and navigators have referred to this phenomenon and tried to explain its

cause. Some of these observations were of commercial value because of the increasing trade and shipping between Peru and Europe via Panama. The Spanish and Portuguese sailors were already aware of the existence of these strong ocean currents. By 1519, they had discovered the advantage of sailing with favorable equatorial currents across the Atlantic to America and returning northward along the Florida Strait in the Gulf Stream as far as Cape Hatteras and then eastward to Spain and Portugal.

Aside from the commercial importance of the currents, there were many pioneering scientists who started to systematically acquire physical and biological information for scientific investigation. In the second half of the XVIII century the thermometer was already available and many navigators wrote of the importance of temperature observations for determination of current boundaries. In 1802, Von Humboldt attempted a global interpretation of the temperature measurements on the Peruvian coastal current [Humboldt, 1926]. Even though he incorrectly concluded that the low temperature was due to the Antarctic origin of the flow (actually the cool water comes from upwelled subsurface water, from Antarctic origin), he was still the first to attempt a comprehensive scientific study and interpretation of this current that still bears his name. After Humboldt, other investigators started to look for additional explanations for this oceanic phenomenon. Because of its importance in the circulation of the Pacific Ocean, many governments, private institutions and entrepeneurs became aware of the benefits of such investigation and began to organize and fund large oceanographic expeditions to acquire scientific data.

B. OCEANOGRAPHIC EXPEDITIONS

In 1823, 1836 and 1837-8, three French expeditions acquired information along the coast of Peru to corroborate Humboldt's theory [Dupery, 1829]. In 1835, the well known expedition by the <u>Beagle</u> accompanied by Charles Darwin visited the coast of Peru and collected biological data [Fitz Roy, 1839]. The assembly of these new data was followed by new theories and interpretations: De Tessan (1844) hypothesized that the low water temperatures were due to upwelling. In 1874, Dinklage added another piece of evidence to the puzzle by pointing out the importance of the trade winds in the westerly set of the surface flow [Gunther, 1936].

The first few years of the XIX century saw a profusion of new theories concerning the explanation of the Peru Current. Sverdrup, Schweigger, Vallaux and Schott were a few of the oceanographers who investigated the Peru Current and the already known "El Nino" countercurrent using more advanced instrumentation and techniques [Gunther, 1936].

During the decade of the 1930's, Sverdrup [Sverdrup, 1930] and Schweigger [Schweigger, 1931] published the first coastal and open ocean observations acquired with subsurface instruments. The data included temperature, salinity, pH and current velocity. These expeditions were so useful in learning about the characteristics of the area that others were encouraged to organize more elaborate data acquisition programs.

In 1928-29, the cruises of the <u>Carnegie</u> [Gunther, 1936] yielded valuable data across the Eastern South Pacific, not only along the coast, but from offshore waters to as far as 115 W and from the Equator to 40 S. Sverdrup's interpretation of the resultant temperature and salinity sections contributed to a better understanding of the limits of the system of oceanic currents and undercurrents along the coast of Peru.

In 1931 the British ship R.R.S. <u>William Scoresby</u> made an extensive survey from the Equator to 40 S acquiring meteorological data and observing temperature, salinity and phosphate concentration from the surface to 400 meters deep. The results of the Scoresby expedition are presented in an extensive report [Gunther, 1936]. After this expedition, and particularly during the 1960's, many well-organized oceanographic cruises have been conducted along the Peruvian coast. The following is a list of some of the most important modern expeditions. Many other short cruises by local and foreign institutions that have taken place in this part of the ocean are not included.

TABLE 1

Data Collection Programs 1950's-1970's

Year	Name	Program or Institution
1952	Shellback Expedition	Scripps Institution of Oceanography
.1953	Yale South American Expedition	Bingham Oceanographic Laboratory
1960	STEP-1 Expedition	Scripps Institution of Oceanography
1967-8	(several)	EASTROPAC Program
1972	El Nino Watch Expedition	Scripps Institution of Oceanography
1976	JOINT II Experiment	CUEA Program

The last three expeditions deserve special commentary because they were large data acquisition and experimental programs that greatly added to the amount of information available from this part of the ocean. Many other cruises done by local institutions also exist and will be incorporated into the bank.

1. EASTROPAC Program

In 1960, at the seventh conference of Eastern Pacific Oceanic Conference (EPOC), the members of this organization discussed the sparse and infrequent data acquired in the Eastern Tropical Pacific. From these discussions, it was concluded that in order to understand and predict the variations of the circulation of this area, a more complete data base was necessary. An ambitious program of geological, geophysical, meteorological, as well as physical, chemical and biological oceanographic data acquisition was proposed. After several years of delay, the original program was somewhat reduced, fisheries surveys were added, and the revised program adopted its definitive name, Eastern Tropical Pacific Program (EASTROPAC). This was to be a cooperative effort towards understanding of the oceanography of the Eastern Tropical Pacific Ocean. The program included the participation of U.S., Mexican and Latin American institutions. The fieldwork was divided into seven 2-month cruises of the area between 20 N and 20 S and from the coast of North and South America to 119 W. Five of these seven periods of observations were conducted adjacent to the Peru coast during the months of: Feb-Mar 1967, Jun-Jul 1967, Aug-Sept 1967, Oct-Nov 1967 and Feb-Mar 1968. These particular cruises included the participation of four U.S. ships and one Peruvian ship. The information acquired was primarily physical and biological oceanographic data but included meteorological observations as well. All the observations during this program were processed and presented in eleven volumes as the "EASTROPAC ATLAS".

EASTROPAC was the first successful international program for the comprehensive acquisition of data in the Eastern Pacific. It was an

instructive experience for all of the participants in the organization of this kind of international cooperative program, and it would prove to be a valuable experience that was to be repeated many more times during the following decade.

2. El Nino Watch Expedition

In October 1974, based on previous studies done on the "El Nino" phenomenon, a prediction was made [Quinn, 1974] that a weak El Nino event would occur in early 1975. A group of scientists sponsored by the Office of Naval Research and the National Science Foundation of the United States organized a multidisciplinary expedition to observe the conditions offshore of Ecuador and Peru. The expedition consisted of two cruises that surveyed the area from 2 N to 14 S, and 95 W to the coast. The first cruise took place from mid-February through March 1975 and the second from mid-April through May of the same year. The data acquired included CTD casts to a depth of 500 m, Niskin casts, XBTs, vertical zooplankton net tows, current measurements and meteorological variables. This information has been analyzed and presented in the "El Nino Watch Atlas" [Patzert, 1978].

3. CUEA Program

In 1966, a group of scientists and government officials in the U.S. proposed a ten year systematic program of cooperative international ocean exploration. This was the beginning of what was to become one of the most significant events in recent oceanographic history, the International Decade for Ocean Exploration (IDOE).

This ambitious program which took place during the decade of the 1970's was motivated by the future uses of the marine resources as well

as by scientific curiosity. It included the participation of many nations. By October 1978, 37 nations were involved in one or more of the various parts of IDOE programs. The original idea of IDOE was to promote an international, interdisciplinary and inter-institutional approach to improve our understanding of the oceans. One of the objectives that was included in the context of IDOE, was that of "Environmental Forecasting". Later on, an objective entitled "Living Resources" was included. Under the auspices of these two objectives, the Coastal Upwelling Ecosystem Analysis (CUEA) program was created. CUEA was a multidisciplinary research program designed to investigate coastal upwelling and its biological consequences. It was planned on the basis of a series of earlier investigations off Oregon and Northwest Africa in the mid-1960's and off the coast of Peru in 1966 and 1969 ∫Costlow and Barber, 1980]. CUEA included several field programs with ships from eight countries participating in the intensive periods of observation. Four small and two large field operations were conducted in the regions of upwelling. The large operations were named JOINT-I, conducted off North West Africa and JOINT-II, off Peru. JOINT-II was a U.S.A.-Peruvian effort of data acquisition and experimentation. It was divided into three periods of intense observations: March-May 1976 (MAM 76), July-November 1976 (JASON 76) and March-May (MAM 77). Seven U.S. and two Peruvian ships participated in the data acquisition. The U.S. provided aircraft, satellites and eight shore-based meteorological stations, and Peru provided logistic support for the aircraft and meteorological stations, as well as a coordination center at the Instituto del Mar del Peru, in Lima.

CUEA has produced one of the most comprehensive sets of data ever collected off the coast of Peru. The results of the field, laboratory and theoretical studies will be a valuable source of data for future investigations in this area.

III. THE IMPORTANCE OF AN OCEANOGRAPHIC DATA BANK IN PERU

A. THE FISHERIES

It is a well known fact that the surface water close to Peru is much colder than what is expected at these latitudes. The temperature was so low that Humboldt thought that the northern current originated in Antarctica. However, in 1844 De Tessan, correctly attributed the low temperature to upwelling of subsurface water (from 100 to 300 m) along the coast. Upwelling is a process whereby subsurface water rises toward the surface. Coastal upwelling results when prevailing winds produce offshore flow in the surface layer near a coastal boundary [0'Brien, 1978].

Areas of coastal upwelling cover only 1% of the oceans, but they contain more than half the world commercial fish stocks. Coastal upwelling occurs along virtually all coasts. It is very intense in several places such as the coasts of California-Oregon, Northwest and Southwest Africa, Arabia, and western South America. In some of these regions, the phenomenon is seasonal (especially in mid-latitudes), whereas off Peru, it occurs throughout most of the year.

Upwelling is of particular importance because it can raise great quantities of phosphates, nitrates, silicates, and other nutrients from subsurface waters. The presence of these nutrients in the photic zone greatly enhances photosynthesis and as a result, algae and phytoplankton flourish. Along the coast of Peru, this great productivity in plant life provides the basis for a prodigious food chain which supports a large fish population, including the largest population of anchovies and

ocean birds found anywhere. As a result, the commercial fishery harvest is enormous. In 1970, Peru accounted for more than one fifth of the total world fish protein production [O'Brien et al, 1981].

However, this productivity at the top of the food chain is not very stable. In the northern part of Peru, a tongue of Equatorial Countercurrent water annually intrudes southward along the coast from the Gulf of Guayaguil. It is of 6 to 7 C warmer and much less saline than the surface water it displaces. This event usually occurs during the Christmas season and is, therefore, known as "El Nino", which means "The Child", the popular name given to the Child Christ, symbol of the Christmas season. "El Nino" occurs every year with its warm waters penetrating to 6° or 7°S. In some years, this countercurrent is stronger than normal and it may reach as far as 12°S or more, causing a catastrophic destruction of plankton and fish life. In "El Nino" years, the large anchovy population which is adapted to the cold upwelled water, either scatters in search of cooler water or it migrates offshore. Many fish die prematurely, dramatically reducing the stock. Large number of seabirds that mainly feed on these anchovies die of starvation, and the guano (seabird fecal deposits used as fertilizer) yield is reduced with a deleterious effect on agriculture. The effects of "El Nino" are evident in the atmosphere as well. In severe "El Nino" years, the northern coastal areas experience torrential rains causing flooding in the farmlands along the river valleys.

The social and economic impact of this event in Peru is well documented [Murphy, 1954; Caviedes, 1975; Paulik, 1981]. In 1960, the fishmeal industry alone represented 35% of total national foreign

earnings [Paulik, 1981]. In 1972, a combination of very intense fishing and a long and intense "El Nino" caused the collapse of the fishing industry. The economically important guano mining industry also felt the effect of this phenomenon in the late 1930's. The economic effects are felt worldwide in the price and supply of fertilizers and fishmeal. In 1973, a sharp rise in the price of chicken in the U.S. was due to the unavailability of fishmeal used as chickenfeed [O'Brien, 1978]. Prices of soybean meal, which competes with fishmeal as animal feed, increased due to reduced supplies of fishmeal on the world market.

The "El Nino" phenomenon is an important example of a complex ocean/ atmosphere interaction whose causes are still not fully understood.

Much effort has recently been made to understand the interaction between atmospheric and ocean conditions that precede the initiation of an "El Nino", yet more data are needed to improve any predictive capabilities for the onset of the event. The far reaching effects of "El Nino" on the Peruvian economy and on the world supply of fertilizers and fishmeal has encouraged the scientific community to try to understand the behavior of this phenomenon by the acquisition of more data. In recent years, there have been numerous studies of this event that used these data. It is now important to update and organize these data to better document "El Nino" and its variability. It is also important to share these data between the countries of the region that are affected by the "El Nino" (Ecuador, Chile, Peru and Colombia) under programs such as ERFEN (Regional Study of the "El Nino" Phenomenon).

B. THE NAVY

The onset of "El Nino" with rapidly changing conditions of the water masses along the coast also changes the acoustical characteristics of the ocean in this area. The Navy is interested in this variability because of its effects on underwater sound propagation for antisubmarine warfare (ASW).

Because of the historical economic impact of "El Nino" and its consequences for Naval operations, it is necessary to develop an oceanographic data bank to archive all available historical ocean data for Peruvian waters and also to update the data bank with new information on a continuing basis to monitor and eventually to predict ocean conditions.

Several years ago, the Peruvian Navy began a study of the physical characteristics of the ocean waters along the coast to provide its ships and submarines with information on the environment in which they operate. The Peruvian Naval Hydrographic Office (DHNM) was assigned the responsibility of collecting data available from local sources. Because of the need for rapid access to the data and for rapid preparation of summaries in different forms, the data were to be in computer accessible form. The DHNM searched for physical oceanographic data in government offices and fisheries institutions but found very little in computer accessible form, and ultimately most of the information acquired came from the U.S. National Oceanographic Data Center (NODC).

This lack of computer accessible oceanographic information from local sources lead to the concept of a data bank for the Navy and other users in which all the data acquired along the coast of Peru for many years by local and foreign sources could be organized in such a way so

as to be of immediate use. The potential benefits from the organization of such a bank include the incentive it will provide for students of marine sciences and other related investigations. Its creation should encourage the Peruvian Navy to implement a strong program of oceanographic data acquisition with Navy ships and other platforms. It should also encourage other sources, local and foreign that acquire oceanographic data to provide it to the bank on an exchange basis. Finally, it will create a mechanism to archive the information gathered by foreign ships that come to Peru on future oceanographic expeditions. On the basis of these considerations, the Peruvian Navy decided to commission the organization of an oceanographic data bank.

C. THE CREATION OF THE BANK

On March 11th 1977, the Government of Peru issued the Decree Law D.S. #033-77-MA by which the responsibility for the creation and organization of the Oceanographic Information System or Sistema de Informacion Oceanografico (SIO), was assigned to the Hydrographic Office of the Navy.

The first step in the organization of such a system is the collection and inventory of all available historical information. As is evident, a large amount of data has been acquired along the coast of Peru for many years. Finding and identifying this data today is difficult because of the diversity of the sources and data formats.

Although a significant effort has been made by various institutions over the years in the acquisition of data, much of this valuable information remains scattered in the archives of those institutions. In recent years, however, efforts have been made to try to centralize and standardize the acquisition of data from all over the world. Two such central

facilities are (NODC) and World Data Centers A and B in Washington D.C. and Moscow, respectively.

The organization of the historical data is only the initial step in the creation of the oceanographic data bank which will support the SIO. Next, there needs to be careful planning for the establishment of procedures to acquire additional data to update the data bank, and for the implementation of such procedures. There are many considerations in planning the future acquisition of data: special projects or areas of study, user requirements, existence of historical data, availability of platforms for data acquisition, Navy priorities, etc.

The creation of the SIO will require that the DHNM organize a network of data-acquisition platforms that can provide the data to update the bank on a continuing basis. This can be done in part by use of Navy ships as well as fishing and commercial ships of opportunity. The use of ships of opportunity has already enabled the monitoring of atmospheric conditions along the coast of Peru, and they could also be used for acquiring oceanographic data. The implementation of such a monitoring system is not simple but can be accomplished by sharing both the responsibilities and the benefits of the program with others. For example, FNOC already has a global ship of opportunity program in existence. Data acquired by these ships provide a large scale monitoring and temperature conditions in the Pacific Ocean and a few of the ships make expendable bathythermograph observations in the Eastern Tropical Pacific.

The implementation of the SIO will be divided into three main stages:


- 1) the acquisition and organization of the existing historical data;
- 2) the establishment of procedures for monitoring, assembling and exchanging of new data to update the data base on a continuing basis; and

3) the implementation of computer analysis programs and application products to provide services for users.

The scope of this thesis is devoted to the first of these steps, that is, the compilation from numerous sources of all available historical data from the coastal waters of Peru. More specifically, the focus is on the design and implementation of an archiving system of primarily physical oceanographic data. One purpose of this thesis is to provide some ideas and examples of possible applications of the available historical data. A final purpose of this thesis is to conduct a pilot study and test of some of the systems already in existence for the management of oceanographic data. The programs developed for the processing of the data will be transferred to the computer facilities of the Peruvian Navy where they will be tested to start the implementation of the SIO. Next, all available physical data will be acquired and introduced to the system to update the bank. Table 2 shows how the flow of information will update the bank in the future.

One of the main accomplishments of this thesis is that it has provided experience in working with programs and data sets similar to those that will be used for the SIO. This experience will be very valuable in the design and implementation of the oceanographic data bank for the Peruvian Navy.

TABLE 2 Flow of Information Into the Bank



IV. ORGANIZATION OF THE BANK

A. THE IMPORTANCE OF DATA FORMATS

The way in which data are recorded is a central issue in the design of any computerized data management system. The computer does not recognize what the data represent; rather, it is the programmer who formats the data for input to the computer as an ordered pattern of binary digits or bits.

There are several different ways of formating oceanographic data. Some formats are more efficient than others depending on the type of data and the applications for which the data are to be used. It is also important that the available computer efficiently supports the format. During the last fifteen years, the improvement of computers and of sophisticated techniques for archiving and retrieving data from large data bases has increased the efficiency of managing oceanographic data. Recently developed systems save considerable amount of memory space and computer time.

As an example of improving data storage techniques, in 1966 the North Atlantic ocean station file of the U.S. Naval Oceanographic Office (NAVOCEANO) was stored on sixty-four reels of tape. By the simple technique of grouping single card images into blocks of ten card images per tape record, the physical size of the file was reduced from sixty-four to twenty-one reels of tape [Yergen, 1970]. This saved considerable space and reduced the complexity of making computer summaries of the data. Later, in the same year, NAVOCEANO restructured the same file

into an experimental format called "rapid access" format Yergen, 1967. This format further reduced the twenty-one reels to three reels of magnetic tape and made the data very accessible.

The new, more efficient scheme used a binary number format rather than the former binary coded decimal (BCD) character format. In this way, the size of the file was reduced and conversions between BCD and binary numbers eliminated. Elimination of the BCD-binary conversion speeds the processing since binary is the natural arithmetic base system of most computers. Reducing the physical size of the file not only saves space and simplifies computer runs but also reduces the actual time needed to transport the tape past the computer's tape-read heads.

There are several disadvantages of binary data, however. Most important is that data are commonly exchanged between computers in character form and many users are unfamiliar with binary data. Also many common compilers (such as IBM FORTRAN) do not contain features to easily handle binary data. Thus, the user is forced to develop special subroutines to pack and unpack the data. Fortunately efficient, FORTRAN callable, machine language subroutines are available for many computers, based on work at the National Center for Atmospheric Research Jenne, 1979 and Scripps Institution of Oceanography Hilton, 1974. These are the routines SBIT/SBITS and GBIT/GBITS which pack and unpack bits or strings of bits.

An example of the volume and computer time saved on a Control Data Computer (CDC) by using binary packed rather than BCD data is given below [Jenne, 1974]. For a number in the range 0 to 4095 in magnitude (such as ocean temperatures in the range -1.8°C to 35.0°C, stored as

hundredths of a degree Celsius, ~180 to 3500 units, and biased by +2.0°C or 200 units to be in the range 20 to 3700 units) the following number of bits are required for storage:

Volume comparison:

one saving in time.

Storage Method	Volume Per Number
Binary Packed	12 Bits
BCD (6 Bit Character)	24 Bits
BCD (8 Bit Character)	32 Bits

Also, the amount of time required to convert the number for use in calculations is as follows:

Type of Data	Time to Unpack It
Binary (12 Bit Number Using Machine Language)	3 microseconds
Binary (12 Bit Number Using Fortran Subroutine)	20 microseconds
BCD Characters Using Fortran Subroutine	120 microseconds
As can be seen, considerable	time and space is saved by using binary
packed data when working wit	h large volumes of data. In the above ex

Ever since the computer became an indispensable tool for storing and processing large amounts of data, many storage devices have been used for archiving and exchanging information between data centers.

Punched cards, punched paper tapes, magnetic tapes, and disks have all been used for this purpose. Today, because of its versatility and low

amples, there is about a three-to-one compaction and about a forty-to-

cost, magnetic tape is a common storage device and its standardization by all major computer manufacturers has simplified the exchange of data on tape between users. However, even though magentic tape has become a standard medium, many different formats are used to store oceanographic data. The increasing diversity of ways of acquiring oceanographic data, together with the variety of applications for the data has encouraged the design of many different formats each suited to the user's particular needs and computer facilities. Based on the earlier work of Yergen, Jenne, and others, the Master Oceanographic Observation Data Set (MOODS) was developed by Fleet Numerical Oceanography Center (FNOC) as an attempt to merge all available ocean profile data globally for rapid access by computers for naval purposes.

B. HISTORICAL DATA FORMAT (MOODS)

The main source of historical data for the coastal waters off Peru was extracted from the MOODS archives of FNDC. During the last few years, efforts have been made at FNOC to acquire all available oceanographic profile data from files of the U.S. National Oceanographic Data Center (NODC) and other sources in order to make the existing oceanographic data easily accessible to operational users. The data conversions and programs to handle the MOODS data were written in Fortran IV by Compass Systems, Inc. and implemented on a CDC 6500 computer at FNOC.

The MOODS system was designed to meet five objectives: (1) simplification of computer software to handle a single format of ocean profile data rather than the multiple formats used previously, (2) compaction of the data files from more than 50 reels of tape to just a few reels (or ultimately, a single disk pack), (3) rapid access by computer, (4) ease

of data exchange between computers of different manufacturers. The data are stored as binary integers, biased to be positive integers to eliminate problems of handling negative binary numbers. For further compaction of the data, values in each profile are stored as differences rather than actual values.

C. DESCRIPTION OF THE MOODS FORMAT

The MOODS files are sorted by time and position (both to the accuracy of minutes), source and ship files in such a way that records can be sorted and extracted in any combination of time, one degree areas, ship and source files. The information packed into a MOODS record, corresponding to one complete ocean profile, is divided into several fields. Each field contains information about the observation in the following sequence:

1. Day-Time Group

This is an eight digit integer that contains the year, month, day and hour of the observation (i.e., 65032110 corresponds to 1000 hours Greenwich time on the 21st of March 1965).

2. Position

Latitude and longitude are stored to the nearest minute. The hemisphere is stored as 2 bits with positive values for north and east hemispheres.

3. Cruise Number and Ship Code

The cruise number usually corresponds to the NODC cruise number. If the data were not obtained from NODC, the number used is the one coded by the source. Because the criteria used by different data acquisition organizations to select their cruise numbers are not the same,

FNOC has developed a cross-reference table to identify the ship codes and names of ships.

4. Number of Levels

This number corresponds to the discrete number of recordings taken during the original observation.

5. Number of Standard Levels

This is the number of levels at which values can be interpolated. A convenient feature of the MOODS subroutines is conversion of the data to standard levels from the original values. The standard depths to which values are interpolated can be selected by the user. This feature is especially useful in mapping the distribution of data at a selected depth, as will be seen in the last chapter.

6. Report Number

All the observations, whether BTs or hydrocasts, are numbered consecutively, and sorted by month, year, and by latitude and longitude (rounded to whole degrees). The report number is generated by a counter that keeps track of the number of observations extracted from the FNOC MOODS files. This number will eventually be altered when installed in the DHNM computer facilities in Peru because additional historical data will be added from other local sources such as the Peruvian Navy and fisheries.

7. Profile Data

The MOODS subroutines contain an array of dimension 7, corresponding to the seven possible variables that can be selected for an observation, coded as follows:

Code 0	Depth
Code 1	Temperature (in degrees Celcius)
Code 2	Salinity (in parts per thousand)
Code 3	Sigma-T
Code 4	Pressure
Code 5	Sound Speed (in meters per second)
Code 6	Oxygen (in millimeters per liter)

More than seven variables may be included in the future (such as chlorophyll), as needed. Any of these parameters or combinations of them can be selected and displayed in a table format, provided the observation contains such information. This feature also enables the user to rapidly select those profiles having the desired parameters.

8. Source Code

The source code identifies the source of the observation and allows the system to sort the data set by source and permits identification of individual observations. FNOC has a table of codes identifying the various sources.

D. EXTRACT FILE FOR THE PERUVIAN AREA

Extraction of all available data from the area off the coast of Peru was done by FNOC from their monthly-sorted MOODS data tapes. The limits of the area selected was from the Equator to 20°S and from 86°W to the coast. This area covers the entire length of the Peruvian coast, approximately 200 miles offshore as can be seen from Figure 1. The extracted information contains NODC mechanical and expendable bathythermographic (BT and XBT) data and hydrocast data, radio messages of BT and XBT data, Scripps Institution of Oceanography file of mechanical BT data read at 100 ft intervals, BT data digitized by Scripps




FIGURE 1: Areas Selected for Data Extraction From MOODS Monthly Tapes

Institution and Compass Systems, Inc., and British XBT and mechanical BT data.

The extracted MOODS data file was read and inventoried at the Naval Postgraduate School (NPS) using the IBM 370 Model 3033 computer. To be able to use the subroutines from the FNOC Control Data Corporation computer to unpack the data in the MOODS format on the NPS computer, the subroutine had to be modified because the Fortran languages used by the CDC and IBM computers are not directly compatible.

This conversion of the FNOC MOODS data read subroutines for use on the NPS computer and will allow the future NPS users to use MOODS format data independent of FNOC's facilities. Now that these subroutines are adapted to the NPS system, it may be worthwhile to install the global MOODS data base at the NPS W.R. Church Computer Center and incorporate the subroutines as part of the public library.

Additionally, since the Peruvian Navy has an IBM computer which will be used for the support of the SIO, MOODS files will be easily accessible using the converted routines by the DHNM for climatological purposes. In addition, new data will be processed by DHNM and will include CTD data acquired on research vessels, XBT data acquired on ships of opportunity operating along the coast of Peru, and sea level data for the six tide gauges installed along the coast. The programs that will process the current oceanographic data for the DHNM will not use the MOODS format but rather a simpler character format because of the elaborate programming required to update MOODS files and the relatively small amount of new data to be processed. The new data will be exchanged with NODC and FNOC and updated MOODS climatological tapes provided to DHNM by FNOC.

E. SELECTION OF THE DATA

Once the MOODS read subroutines were converted to the IBM computer and tested, the data were separated into two groups: bathythermographic (mechanical BT, XBT, or AXBT) and hydrocast data. The data were then listed in a convenient format (examples are shown in Appendix A). In the following chapter, it will be seen that the separation of data into hydrocast stations and BTs was done to form independent sets of data for specific applications. For example, sound speed profile analysis requires salinity information and thus can only use hydrocast data.

V. APPLICATIONS

A. INTRODUCTION

The two primary functions of the SIO are: archival of data concerning the ocean area off the coast of Peru and distribution of this information to the users. The applications of the data are the most important aspect in the consideration of the design and organization of the data bank. The effectiveness of any information system depends more on how well it satisfies the users' needs than on the level of sophistication of the system.

It is of vital importance to clearly define the problems to be solved by the users to best define the organization of the information system. Based upon the priorities and objectives of the system, the design and organization of the data bank can be suitably tailored. Many of the potential users of the SIO and their specific needs have already been identified. These are the Navy, the fisheries, other agencies and scientific researchers. In addition, there may be other potential users having specific requirements for the use of this system in Peru. Many of these are difficult to identify at this time, and their presence will be obvious only after the system is created. It is for these cases that the flexibility of the system becomes important so that it can best satisfy the demands of all users with a minimum of data processing.

Access to the SIO probably will initially be limited to a number of users having very different specific needs. However, it is very likely that additional users having other requirements may access the system

later on. Such new requests will encourage the growth and diversification of the system as well as provide new ideas for the acquisition and utilization of other oceanographic variables and the implementation of new applications.

This thesis primarily dealt with physical oceanographic data because very little of these data exist in the archives of any Peruvian institution in comparison to the large amount of chemical and biological information collected by the Instituto del Mar del Peru, the Ministry of Fisheries and other sources. Many of these biological and chemical data will eventually also become part of the data bank of the SIO.

In the following sections some of the more immediate applications and products that can be derived from the historical data collected for the data bank are presented, and other ideas of probable future use of these data are discussed.

B. INVENTORY OF THE DATA

The first task in organizing the bank is conducting an inventory of the data.

The criterion used in selecting the variable under which the data are filed is very important, and it will also depend on the specific needs of the user. The system has to be flexible enough so that different sorting parameters can be used to select the data to be retrieved and presented to the user. For purposes of illustration, two cases of retrieving the data by different parameters will be shown. These are inventories of the observations by years and by months.

The selection of year as a parameter to retrieve and display the information is useful, and it is a familiar way of displaying data sequentially. Data presented this way is particularly useful when studying phenomena of very low frequency of occurrence; such as, "El Nino".

As stated earlier, the information collected includes bathythermographic (BT) and hydrocast observations. In order to provide a graphical representation of the geographical distribution of these two kinds of observations, three different maps were made: BT data, hydrocast data, and both types of data (Figures 2, 3 and 4). To complement the maps of the distribution of the data, a listing with the basic information of the individual observations is provided in Appendix B. The date, source and time of each observation can be obtained from this listing.

A second way in which the information can be retrieved and presented is by month. This is useful in the investigation of phenomena having regular seasonal trends. One example of the application of monthly data is the selection of representative sound speed profiles by seasons (quarterly) for certain geographic areas. This application will be discussed in more detail in Chapter IV. The monthly observations are displayed in three maps similar to that for the yearly data (Figures 5, 6 and 7). Listings for this example are also given in Appendix B.

These were just two of the many possible ways in which data can be selected and displayed. Other applications that can be derived from these data could be tables of the distribution of the data by source, year or other parameter. There can be multiple ways of selecting and displaying the data, and the user ultimately will decide the way the data are presented and sorted for specific applications.




FIGURE 2: Distribution of BT Historical Data for 1960




FIGURE 3: Distribution of Hydrocast Historical Data for 1960




FIGURE 4: Distribution of Hydrocast and BT Data for 1960




FIGURE 5: BT Historical Data for the Month of June, All Years




FIGURE 6: Hydrocast Historical Data for the Month of June, All Years




FIGURE 7: Hydrocast and Historical BT Data for the Month of June, All Years

C. NAVAL APPLICATIONS

Two of the purposes of the organization of the SIO are to provide the Peruvian Navy the capability of obtaining oceanographic data for analysis and to provide operational products for naval applications. Some of the practical applications derived from the processing of historical oceanographic information for the Navy have been tested for several years, and some of these have been implemented as operational products in many navies. Some examples of these products are those used in navigation aids, environmental forecasts, and acoustic analyses. In this section, one example of an application for the Navy using the SIO that can readily provide a useful operational product will be given. The discussion of this example is the subject of another NPS thesis [Garcia, 1981] that used data selected directly from this bank.

The acoustical characteristics of the water masses in a naval operation area are of vital importance when planning and executing submarine and ASW operations. Although the fleet normally acquires BT's during operations to analyze the thermal structure of the ocean, information is needed prior to the operation for planning purposes. Historical data can be used to analyze the ocean conditions statistically by geographical quadrangles and time of the year.

To determine the acoustic characteristics of the water masses along the coast of Peru, the geographical area between 0° and 20°S and 70° to 86°W was divided into 83° one-degree quadrangles (Figure 8). The details of the selection of the areas as well as the analysis of the data for each area is given in Garcia (1981). As an example, an analysis of one of the sections will be summarized here. To study the acoustical





FIGURE 8: Map with Grid of One Degree Quadrangles

effects of a water mass in each of the areas, the available historical data were used to obtain monthly graphs of sound speed profiles. These profiles were grouped by quarters: Jan-Mar, Apr-Jun, Jul-Sep and Oct-Dec. They were then plotted separately for each one-degree quadrangle and quarter. (Figure 9 represents all the profiles from quadrangle #60 for Apr-Jun). From each of the quarterly groups of profiles, one or more curves were selected as the ones that best represented the characteristic of the water mass for that area and quarter. This selected profile was then input to an acoustical model to obtain curves of transmission loss versus range. Figure 10 shows the transmission loss curves for 6 frequencies (50, 100, 200, 300, 500 and 800 hertz) using the representative sound speed profile for April-June for quadrangle #60 of Figure 9. In this particular example, the acoustic model used is the Fast Asymptotic Coherent Transmission Model (FACT), implemented at the W. R. Church Computer Center at the NPS. The transmission loss curves from the FACT model provides detection ranges for specific sonar characteristics of a given ship. This is strategically vital information when planning naval operations involving submarines and ASW. Other kinds of information that can be readily obtained and also useful for ASW applications include surface temperature charts and charts of the average depth of the thermocline.

D. OTHER APPLICATIONS

1. Topography of Isothermal Surfaces

In recent years, many new theories and models have been developed to predict and simulate the onset of "El Nino" [Hurlburt et al., 1976; Wyrtki, 1975; Bjerknes, 1961]. The data acquired by the CUEA experiments



___ FIGURE 9: Sound Speed Profiles for Period April-June Quadrangle #60




FIGURE 10: Trans. Loss Curves for Representative Profile at 6 Frequencies

and others in the Pacific during the last decade have been analyzed in light of the new theories with very interesting results. Wurtki (1975) suggested an explanation for "El Nino": strong southeast trade winds over the Eastern Pacific produce an east-west slope of sea level and an accumulation of water in the western Pacific. When the south-east trade winds relax, the accumulated water moves eastward as an equatorial internal Kelvin wave. Wyrtki found good correlation between "El Nino" years and weakening of the south-east trade winds in the previous year. As a result of the weakening of the trade winds, warm water from the west flows toward the northern coast of Peru, depressing of the thermocline. The effect of the depression of the thermocline can be clearly seen by mapping the topography of an isothermal surface. The 15°C isotherm in the area off Peru lies in the thermocline and intersects the sea surface during strong upwelling in the Southern winter. Thus, the 15°C isotherm is a good indicator of the depth of the thermocline and can be used to map areas of upwelling. Based on published cruise reports, Wyrtki (1975) presented maps of the topography of the 15°C isotherm surface during several months in 1972 and showed progressive deepening of the surface as a result of the strong "El Nino" in that year.

Climatological data can be used to map topography of an isothermal surface and, thus, to determine the variability of the depth of the thermocline. Figure 11 shows the depth of the 15°C isotherm computed from the SIO for data acquired in May 1971. If sufficient data are available, a sequence of maps can be obtained to show the variation of the depth of the thermocline and of areas of upwelling.




FIGURE 11: Map of Depth(m) of 15 C Isotherm for May 1971

The fluctuation of the depth of other isothermal surfaces have also been analyzed for similar studies. For example, the annual variation of the slope of the 14°C isotherm along the equator has been correlated with wind stress values to study the eastward propagation of energy in this region [Meyers, 1979].

2. Sea Level Data

Sea level data, as recorded by the DHNM at six tide gauge stations along the coast of Peru, will be part of the SIO as shown in Table 2. The recording and study of sea level and its fluctuations have been done for centuries. The ancient Greeks were the first to record the relationship between the tides and the moon's monthly passage around the earth. It was not until Newton presented his gravitational theory that this relationship was explained. Since then, many different techniques have been used for recording and predicting the sea level.

Because of its importance to navigation and its ease of measurement, sea level is now widely recorded. Historical sea level data exists for periods of decades or longer at many coastal and island locations worldwide. Sea level fluctuations have been closely studied recently, to study low frequency fluctuations resulting from various oceanic and atmospheric processes.

Along the coast of Peru, several studies have recently correlated winds and surface temperature changes with sea level changes. The results indicated the presence of low frequency perturbations traveling poleward as coastally trapped baroclinic waves (internal Kelvin waves) [Smith, 1978; Enfield and Allen, 1970]. These waves seem to play an important role in the propagation of an "El Nino".

VI. CONCLUSIONS

Large amounts of data have been acquired along the coast of Peru for many years. Some of these data have already been assembled by large oceanographic data centers like NODC, World Data Center, and Fleet Numerical Oceanography Center. However, there are still some historical data that have not been incorporated in those data banks and can be found only in the archives of the original sources.

A large proportion of the oceanographic data acquired along the Peruvian coast, particularly during the last decade, was obtained under international cooperative programs and experiments. Most of the information obtained by the international programs and by Peruvian Institutions is generally available. However, no Peruvian agency or institution was assigned the responsibility of assembling all available data acquired during those programs.

In order to improve the understanding of the effects of currents along the coast of Peru as well as the "El Nino" event, it is important to organize an oceanographic data bank in Peru. Most of the historical data required for such a bank are available from large international oceanographic data centers and from Peruvian and foreign institutions. Unfortunately, some of the latter information is not in a computer accessible form. Also, some of the data that are in computer accessible form are stored in non-standard formats, making the merging of all the available data difficult.

The Peruvian Navy, as one of the institutions that acquires oceanographic data on a continuing basis, was recently assigned the responsibility of organizing an oceanographic data bank for Peru. The first step
in the organization of the data bank was to compile, from several sources,
all the readily available historical data from the coastal waters of
Peru. The next and most important step will be the establishment of
procedures to update the data bank on a continuing basis. This will involve monitoring ocean conditions and exchanging data with other institutions. This program will have to be done in cooperation with local
institutions such as fisheries as well as regional and international
organizations under a data exchange agreement. The last step in the
organization of the data bank will fulfill its mission: the implementation of computer programs and application products to provide a variety
of data analysis services to users.

APPENDIX A

FORMAT OF HYDROCAST PROFILE

REPORTE NRO 613	GRUPO FECHA HORA = BUQUE = 00061EX TIPO	65012116
COCIGO DE LLAMADA DEL	BUQUE = 00061EX TIPO	DE DATA= 4
POSICION : LAT =-	-18.50 LONG =-72.38	

NIVELES = 13	TEMP.	SAL.	VEL.SON	OXG.
G.00 10.00 20.00 30.00 50.00 74.00 995.00 1997.00 1997.00 496.00	25.01 24.77 20.20 15.68 13.95 11.877 8.14	34.93 35.99 35.99 35.99 33.33 34.99 34.99 34.66 34.66	1534.80 1534.80 15527.50 15509.20 1509.10 1509.30 15097.90 1495.20 1491.20 1486.60	5.40 5.00 6.020 6.250 7.70 0.370 0.30 0.30 0.30
992.00	4.87	34.61	1486.30	1.50

FORMAT OF BT PROFILES

REPORTE NRO 6592 GRUPO FECHA HORA = 62063006 CODIGC DE LLAMADA DEL BUQUE = 14524 TIPO DE DATA= 3 PCSICION: LAT =-12.78 LONG =-77.90 NIVELES = 11

NIVELES =	TEMP.	PROF.	TEMP.	PROF.	TEMP.
0.0	27.03	10.00	25.73	20.00	21.97
30.00	17.65	55.00	15.72	70.00	13.31
95.00	11.07	180.00	10.05	220.00	8.40

APPENDIX B

LIST OF OBSERVATIONS - YEAR 1960

******	*****	ARCHIVO A	*********** NO 1960 ******	*****	****	****
NRO	GR.FECHA	CDD.aUQ	LAT	LONG	T	++ ++ H
345677890167890101234123457890123456666666667777777777777777777777777777	118 600121122001 221122001 221122001 221122001 2212220077899565801 600012222222005801 6000122222222005 60000122222222005 60000122222222000 60000122222222000 60000122222222000 60000122222222000 60000122222222000 60000122222222000 6000012222222000 6000012222222000 6000012222222000 60000012222222000 60000012222222000 60000000000000000 60000000000	33333333333333333333333333333333333333	0832875332008720057830875332322200002755203352857 12689135795568139324566666	8742198654998621356490012290001123345677888890111 8742198654998621356490012290001123345677888890111 88888888888888777777777777777777777	๛๚๚๛๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚๚	11222 1 1 1 2 3 3 3 3 1 3 3 3 3 3 3 3 3

45 67893456789012347890123128945232345678934567890123456456734567 3333334444444555555555666699003388844444333333111111334444444111133333 3433333344444444	900011122133667778230001112231456702331112233499902333001211211122133663000111221112213366300011122111122300000011122111211122300000011122111122111122000000111221111221111222000000	I I I I I I I I I I I I I I I I I I I	58258255523825250377038870700828035888802222538889250308875037880505 	88888880003555783020855550082073377222273530757520778587530883333333333333333333333333333	๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛	3 3 3 3 3 3 3 5 1 9001123936677823563001125909978301301238901234567023112349902312233 11111111111122 2 111 2 22 1122211111222 111
---	--	---------------------------------------	--	---	---	--

```
33333333433
                                                                                                                                                     601288995
5
                                                                                                                                                  1112233689922770011455663393313022561212491000382223004848
11121111111122 1 3343 3 355341 2 3343 3
```

10891 60100310 13406 HO 10902 60100201 13406 10902 60100302 13406 10903 60100302 13406 HO 10904 60100303 13406 HO 10906 60100305 13406 HO 10908 60100308 13406 HO 10910 60100309 13406 HO 10911 60100309 13406 HO 10912 60100308 13406 HO 10913 60100308 13406 HO 10914 60100309 13406 HO 10937 60100417 13406 HO 10938 60100318 13406 HO 10938 60100312 13406 HO 10938 60100312 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100314 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100314 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100314 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100312 13406 HO 10988 60100314 13406 HO 10988 60100312 13406 HO 10988 60100314 13406 HO 10988 60100312 13406 HO 10988 60100314 13406 HO 10988 60100407 13406 HO 11044 60100407 13406 HO 11044 60100407 13406 HO 11044 60100409 13406 HO 11045 60100409 13406 HO 11046 60100409 13406 HO 11047 60100409 13406 HO 11049 60100409 13406 HO	227728833222883300508205883557050725000037888825730588752878088811	1 3 51 24 20031122356689678013302345224446789C123901212378901012423401567892234545 11 11 11 11 11 11 11 11 11 12 12 11 11 1
---	--	---

11099 111223 601006 601006 111224 601006 111225 601006 111226 601006 111227 601006 601006 111227 601006 601006 111226 601006 601006 111227 601006 601006 601006 601006 111228 601006	13440666 HO 13740666 HO 1374066 HO 13740666 HO 13740666 HO 1374066 HO 137406 HO 1374	05333323388002222233875320330075323350382337770228508878005578077555879000123335677890001234575598887689245066544322119887556666666666666677777777777777777777	20000733000035503330505050038385055580387778858203773327233858227008 3333357789912446799013467889902357779146888664431057410897655320003220 311111111111122222223333333333	๛๛๛๛๚๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛	7802301223567856891230122301234578023124478812370C12645667901228993 11211222222 111121111222 11 111211222 11 11
---	--	--	--	---	--

115223 60100616 039222 134066 115223 601002622 133406 6 115227 601002622 133406 6 115227 601002700 133406 6 1155227 601002700 133406 6 1155227 601002700 133406 6 1155229 601002700 133406 6 1155230 601002700 133406 6 1155230 601002700 133406 6 1155384 60101704 133406 6 115588 60101706 133406 6 115589 60101700 133400 6 6 115589 60101602707 133400 6 6 115599 6010027707 133400 6 6 115599 6010027707 133400 6 6 115599 6010027707 133400 6 6 115599 6010027707 133400 6 6 115599 6010027707 133400 6 6 115599 6010027708 133400 6 6 115599 6010027708 133400 6 6 115599 6010027709 133400 6 6 115599 6010027709 133400 6 6 115599 6010027709 133400 6 6 115599 6010027707 133400 6 6 115599 6010027707 133400 6 6 115599 6010027707 133400 6 6 115600 6010027707 134400 6 6 115600 6010027707 134400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 6010027709 133400 6 6 115600 601002700 133400 6 6 115600 601002700 133400 6 6 116600 601002700 133	3070370358535720322502582358302388385285283855732735500882225777553 1402346789321086544967235677791311187640235688954210077012234566789 2422222222222222333333333333333334444444	05702358037372852330238257080270330030857803553507235880722255033285 41665432219865753110969987767676855475305433211397420033567780233568 7777777777711122223666666666667777777888888887777777777	๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๗๑๗๗๗๗๗๗๗
--	--	---	---

11761 60102922 08761H0 11823 60103001 13406 11824 60103003 13406 11825 60103003 13406 11827 60103006 13406 11828 60103007 08761H0 11829 60103012 13406 11832 60103012 13406 11832 60103012 13406 11833 60103013 13406 11833 60103014 13406 11833 60103015 13406 11833 60103015 13406 11839 60103015 13406 11891 60103017 13406 11891 60103017 13406 11891 60103018 08761H0 11892 60103018 08761H0 11893 60103018 13406 11894 60103017 13406 11895 60103100 13406 11896 60103101 13406 11897 60103101 13406 11896 60103101 13406 11897 60103101 13406 11897 60103101 13406 11928 60103101 13406 11939 60103101 13406 11930 60103101 13406 11944 60103102 13406 11931 60103114 13406 11932 6010312 13406 11933 6010312 13406 11934 6010312 13406 11935 6010312 13406 11931 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11934 6010312 13406 11931 6010312 13406 11932 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11931 6010312 13406 11932 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11933 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11933 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11933 6010312 13406 11931 6010312 13406 11932 6010312 13406 11933 6010312 13406 11931 6010312 13406 11932 6010312 13406 11932 6010312 13406 11933 6010312 13406 11933 601031 14 13406 11931 601031 15 13406 11932 601031 10 13406 11932 601031 10 13406 11933 601031 10 13406 11931 601031 10 13406 11932 601031 10 13406 11933 601031 10 13406 11933 601031 10 13406 11931 601031 10 13406 11932 601031 10 13406 11933 601031 10 13406 11933 601031 10 13406 11934 601031 10 13406 11931 601031 10 13406 11932 601031 10 13406 11932 601031 10 13406 11933 601031 10 13406 11931 601031 10 13406 11931 601031 10 13406 11932 601031 10 13406 11932 601031 10 13406 11933 601031 10 13406 11931 601031 10 13406 11931 601031 10 13406 11931 601031 10 13406 11931 601031 10 13406	355333557802050588035053000388802772828350570703507088707888882200772901234445689966011345789011246890029942063108642100863197531100996456666666677777777777788888888899222223333444444445555566666677777777777788888888992222233334444444555556666667777777777788888888899222223333444444455555666666777777777788888888899222223333444444445555566666677777777778888888888992222233334444444455555666666777777777788888888899222223333444444445555566666677777777778888888888992222233334444444445555566666677777777778888888888	525837333702358555238038733752000038580288888875558252578280880830780124566780188024468035579914791334444554444444444444449022332344667788888889994569999900400011112221111111111111111111111	<u></u> <u>๎</u> <u></u> ๔๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓	433333 43333 33 3 3 32 3 3 2 4 4 2 5 34 42 212345672345672345012300115678 212345672345012300115678 212345672345012300115678
--	---	---	---	---

12597 12794 60112311 03869 12795 60112312 03869 12797 60112313 03869 12798 60112315 03869 12930 60112305 03869 12931 60112307 03869 12932 60112308 03869 12932 60112308 03869 13003 60112308 03869 13003 60112308 03869 131308 13004 60112308 03869 131318 60112300 03869 131318 60112300 03869 13141 60112300 03869 131439 60112220 03869 131440 60112220 03869 131440 60112220 03869 131441 60112221 03869 131440 60112301 03869 131440 60112302 03869 131440 60112303 03869 131440 60112304 03869 131440 60112304 03869 134066 110702 134066 110702 134066 110702 134066 114003 13406 60110007 134066 14003 14004 60110007 13406 14003 14004 60110007 13406 14003 14004 60110061 13406 14037 60110610 13406 14037 60110601 13406 14038 60110610 13406 14037 60110601 13406 14038 60110610 13406 14037 60110600 13406 14038 60110610 13406 14126 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110610 13406 14131 60110511 13406 14161 60110101 13406 14177 60110102 13406 14177 60110104 13406 14177 60110108 13406 14177 60110108 13406 14177 60110108 13406 14177 60110108 13406 14177 60110108 13406 14177 60110108 13406 14177 60110108 13406 14177 14181	07735330078377832527003770038308550877328700025008520507380 27531086411008764203316552213699989872099764333186664322356890 78888999999900000111777777777777777778888889999999999	528572387878558022300877722227057877838200527555385008508003552 81123478992108934562459883321103653109888542100975553214791350 800000000000000000000000000000000000	๛๛ 4 ๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛	91234556798893012340121229057800567303411789001067881231245784
---	--	---	--	--

LISTING OF OBSERVATIONS - MONTH OF JUNE

*****	********	**** **** ** MES DE	JUNIO	****	****	****
NRO	GR. FECHA	COD BUQ	LAT	LON	*****	НО
* *N-66666666666666666666666666666666666	GR.FECHA 61060717 61060717 61060719 61060719 61060723 61060723 61060803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 610608803 661060803 661060803 666060803 666060803 6660609903 6660609903 6660609903 6660609903 666060903 666060903 666060903 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803 666060803	**************************************	*******	LON 33500773003007730077300773007777777.6.68800113555800077300773077777777788.6.5580007300773077373377300307725788.6.88111.6.6360055561.611755768.6.88111.6.88112.2.2.3.3.3.6.8817.0.88111.6.88112.2.2.3.3.3.6.8817.0.88111.6.88112.2.2.3.3.3.6.8817.0.88111.6.88112.2.2.3.3.3.6.8817.0.88111.6.88112.2.2.3.3.3.6.8817.0.88111.6.88111	* *	* *H-11112222 11111222 11121122 1 112112 11 11
239012334567012890123453775566666666666666666666666666666666	5660612215 6660612215 6660612215 6660612215 6660622416 666062245013 666062245013 6660622600 66606226018 66606262618 666062628111 666062628111 6660629113 6660629113 66906181	10111111111111111111111111111111111111	88077735500888225522558890 	-78.08 -78.08 -78.08 -78.08 -78.08 -79.08 -77.09	าตตลงสตรสตรสตรสตรสตรสตร	1527573563034128980121374 112112 11 1112111111

5678456789045656666666666666666666666666666666666	\$344502913133270024790148234256802445793502303813135802131469170111122122002002011122011122000011122000111122000011122000000	N N N N N N N N N N N N N N N N N N N	52052200333777775878888377880308388380883858230373008530238880728998333333333333344444444444555555555555	302305880008777708088088882522502372827333323383822875258785238053203023058800087777777777777777777777777777777	ADMANAMANAMAAAAAAAAAAAAAAAAAAAAAAAAAAAA
67661 677663 6777665 67776667 677777777777777	69060412 69060423 690605004 690605004 69060501 690605117 690605117 69060521 69060520 69060601 69060601 69060604	11111111111111111111111111111111111111	-15.75 -15.6338 -15.63488 -15.63727 -15.633727 -15.63827 -15.63827	525878523805320 -777777777777777777777777777777777777	

The state of the s

56666666666666666666666666666666666666	690606112 690606112 690606112 690606114 690606116 690606116 690606116 6906061113 690606077013 6906077113 690607113 690607113 690607113 690607113 690607113 690607113 690607113 690607113 69060713	N NNN N N NNN N NNN N NNN N NNN N NNN N	32730285027582825887358573873888 23123212321222132222361635135588 5555555555555555555555555555555	53832083200338028802802802202888 	๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๓๚๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛
789012345678901234567890234123456123478901234567890167801123456780 777788888888899999999999999999999999	690662201011110013627903938057134679166622012035168116013362790393805713467916662201203516816013303111112200122122122003571346790666221212121221221221221221220035713467912000666199066119111111122000661191101111111111	15092 15092 00381SN 00381SN 15051 15092	32730285027582825887358573873888700888033333207037275557738587757723123212321222132222361635135588733888700888033333207037275557738587757723123212321222132222361635135588733767712343323648579930044244624665555555555555555555555555555	538320832003380288088028022028885337830752787532025355305770772076 34334533454344444432454795777777777777777777777777777777777	๚๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛๛
6858 6860	69061921	00381SN 15092	-16.67 -16.58	-74.17 -74.32 -75.18	3

1289990167892789011967890123678901223901223456789012234567890122345678999999999999999999999999999999999999	02440093103153713501900366049230253813562355567883802179235066622166128022179235206622161811922000001181111111111111111111111111	N N N N N N N N N N N N N N N N N N N	2822207078783050850235330088835225585250302235333373058588550 1533498904040585387867323311961664488744211122334469862247005 666667777777788888889999999999999990000000000	0855302035357757752080533388387885570500773377557753752222827	ՠ ቀՠቀՠՠ֍֍ՠՠ֍֍ՠՠ֍֍֍֍֍֍ՠ֍֍ՠՠ֍֍ՠ֍֍ՠ֍֍ՠ֍֍ՠ֍֍ՠ֍
26781222906012 255555588906012 2555555566666666666666666666666666666	70062218 70062222 70062301 70062307 70062609 70062303 70062303 70062303 71060417 71060423 71060423 71060217 71060218	133444444 1511333344444 1551133333333333	985858855508860727 	-77.72 -77.922 -777.922 -778.062 -80.633 -81.337 -81.42 -81.40	, חחחחח4ח4חחח 4

6303 71060222 005519 -4.00 -811.33 6304 71060508 151669 -4.00 -82.20 6316 63107 -4.000 -82.20 63108 -4.000 -82.20 63108 -4.000 -82.20 6311 71060510 151569 -4.000 -831.33 6311 71060510 151569 -4.000 -833.33 6311 71060510 151569 -4.000 -833.33 6311 71060510 151569 -4.000 -833.33 6311 71060510 151569 -4.000 -833.33 6311 71060510 151569 -4.000 -831.33 6311 71060510 151569 -5.048 -8811.33 6311 71060200 151569 -5.048 -8811.33 6311 71060200 151569 -5.048 -8811.33 6311 71060200 151569 -5.048 -8811.33 6311 71060201 151569 -5.048 -8811.33 6314 71060201 151569 -5.048 -8811.33 6316 6317 71060211 005510 -5.048 -8811.33 6316 6317 71060211 005510 -5.048 -8811.33 6316 6317 71060211 005510 -5.048 -8811.33 6316 6317 71060211 005510 -5.048 -8811.33 6316 6317 71060211 151699 -6.0755 -8811.33 6316 63324 710606011 151699 -6.0755 -8811.33 63324 710600201 151699 -6.0755 -8811.33 63344 710600207 151699 -7.755 -8800.97 63346 710600207 151699 -7.755 -8800.97 63346 710600614 151699 -7.755 -8800.97 63346 710600207 151699 -7.755 -8800.97 63346 710600207 151699 -7.755 -8800.97 63360 -8800.97 63600 -8800.97 63000 -8	33434343434343434343434333340000000000	2 1111 112 11 122 11 12 11 2 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 12 11 11
--	--	--

589502378925125684589078078905238901340367834073024912208993333333333333333333333333333333333	72062614 72062323 72062620 72062605 72062615 72062702 74061200 74061215 74061215 74061317 74061117	15194 00651UN 15194 15194 00651UN	-17.67 -15.73 -17.67 -17.67 -17.65 -17.63 -4.25	-73.50 -74.70	949944444444	122 1 121 1 2 111212 11 1 2121 12 1 1 1 1 1 2121 12 1
6729	72062620	15194	-17.67	74256832753933192837248539934822935833272445	ġ	ŽÕ
6805	72062702	15194 00651UN	-17.67 -17.65	-75.27 -72.57	4	5
6892	72062615	00651UUNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	-17.67	-73.60	4	15
6903	72062702	00651UN	-17.63	-75.80	4	2
6308	74061206	00721SN	-4 .25	-75.80 -81.35 -81.28 -81.70	4	6
6305	74061209	00721SN	-4.25	-81.70	4	. 9
6315	74061215	00/215N	-4.25 -4.25	-82.52 -83.30	4	20
6321	74061117	007215N	-5.18	-81.92	4	17
6322	74061309	00721SN	-5.12	-81.32	4444444444	. 9
6326	74061109	00721SN	-5.33	-83.17	74	14
6328	74061109 74060921 74061000 74061003	00721SN	-6.95	-79.97	4	21
6334	74061000	00721SN	-6.95 -6.95	-80.28	4	Q
6348	74061009 74061014	00721SN	-6.12	-81.32	4	9
5349	74061014	00721SN	-6.28	-81.70	4	14
6357	74061418 74061019	00721SN	-6.95 -6.62	-81.22 -82.45	4	19
6358	74061023	007215N	-6.80	-82.83	4	ŽŹ
6370	74060814 74060820	NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	-7.77 -7.79	5302257778020253752008 -8311	4	14
6378	74060920	00721SN	-7.75	-80.92	4	20
6379	74060900 74060915 74060918 74060904	00721SN	-7.42	-80.90	4	15
6380	74060918 74060904	00721SN	-7·12	-80.30 -81.48	4 4 4	18
6392	74460704	00721SN	-8.78	-78.83	4	4
6393	74060707	00721SN	-8.78	-78.27	4444444	, 7
6409	74060715 74060803	00721SN	-8.48	-80.45	7	73
6410	74060803 74060807 74060723 74060618	00721SN	-8. <u>20</u>	-80.38	4	7
6411	74060723	00721SN	-8.77 -9.53	-81.50 -79.97	4	23
6454	74060622 74060613	007215N 007215N 007215N 007215N 007215N 007215N 007215N 007215N	-9.15	-78.837 -78.2255 -80.9357 -80.583 -819.3327 -79.338 -78.727 -78.777 -78.277	4	22
6460	74060613	00721SN 00721SN	-9.92	-80.38	44444	13
6556	74060600 74060518	007213N	-10.13	-78.75	4	18
6617	74060420	00721SN 00721SN	-12.07	-77.27	4	ŽŎ
6618	74060500 74060504	007215N	-12.03	-77.77 -79.29	4	Ŏ
6644	74060508	00721SN	-12.05	-78.28 -78.78	4	8
6650	74060508 74060512 75060200 75062906 75060112	007215N 007215N 007215N 007215N 007215N R160	30555582535552852078522088888073523873357733378 	-77.77 -78.28 -78.78 -79.28 -85.93 -85.92 -84.13 -83.37 -81.00	4 4 4 5 10 5 5 10 10	12
6363	75062200 75062906	R160 32U4TM R160 R160 R160 32U4TM 32U4TM	-5.// -6.83	-85.93 -85.90	10	9
6390	75060112	Ř 160	-7.13	-85.90 -84.12	~ <u>š</u>	12
6412	75060100 75062918 75063006	R160	-8.53	-82.33 -83.37 -81.00	15	,0
6499	75063006	32U4TM	-10.68	-83.37 -81.00	ίŏ	7.6

LIST OF REFERENCES

- Bretschneider, D. E., 1980: Sea Level Variations at Monterey, California, Master Thesis, Naval Postgraduate School, Monterey, CA.
- Bjerknes, J., 1961: "El Nino" Study Based on Analysis of Ocean Surface Temperatures, 1935-57, Inter-Amer. Trop. Iuna Comm. Bull., 5, 219-307.
- Caviedes, C., 1975: El Nino 1972: Its Climatic, Ecological, Human and Ocean Implications, Geograph. Rev. 65, 493-509.
- Costlow, J. D., and Barber, R., 1980: IDOE Biology Programs, Oceanus, 23 (1), 52-56.
- CUEA Data Reports (67 Vols.) and CUEA Technical Reports (61 Vols.), National Oceanographic Data Center, Washington, D.C.
- Duperey, L. I., 1829: Voyage Autour du Monde Sur la Corvette de la Majeste, "La Coquille" Pendant les Annes 1822, 1823, et 1825, <u>Hydrographie et Physique</u>, Paris.
- Enfield, D., and Allen, J. S., 1980: On the Structure and Dynamics of Monthly Mean Sea Level Anomalies along the Pacific Coast of North and South America, J. Phys. Oceanogr., 10 (4), 557-578.
- Fitz-Roy, R., 1839: Narrative of the Surveying Voyages of his Majesty Ships Adventure and Beagle between the Years 1826-36, II, 505.
- Geskell, T. E., 1973: The Gulf Stream, The John Day Company, New York.
- Glantz, M., and Thompson, D., 1981: Resource Management and Environmental Uncertainty, Advances in Environmental Sciences and Technology, Wiley Interscience, 11, 3-13.
- Gunther, J. L., 1936: Report on Oceanographic Investigations in the Peru Coastal Currents, <u>Discovery Rep.</u>, 13: 107-276.
- Humboldt, A. Von, 1822: <u>Personal Narrative of Travels to the Equinoctial</u> Regions of the New Continent during the Years 1799-1804 (Translated by Helen Maria Williams), London, 1822-29.
- Hurlburst, H. E., et al., 1976: A Numerical Simulation of the Onset of "El Nino", J. Phys. Oceanogr., 6, 621-631.
- Jenne, R. L., and Joseph, D. H., 1974: Techniques for the Processing Storage and Exchange of Data, National Center for Atmospheric Research, Tech. Note NCAR-TN/IA-93.

Murphy, R., 1937: Notes on the Findings of the William Scoresby in the Peru Coastal Current, Geograph. Rev., 27, (2).

Murphy, R., 1954: The Guano and Anchoveta Fisheries, Resource Management and Environmental Uncertainty, Wiley Interscience, 11:81-106.

O'Brien, J., 1978: El Nino: An Example of Ocean-Atmosphere Interaction, Oceanus, 23, (4):40-46.

O'Brien, J., et al., 1981: Ocean Models of "El Nino", Resource Management and Environmental Uncertainty, Wiley Interscience, 11.

Paulik, G., 1971: Anchovies, Birds and Fishermen in the Peru Current, Environment: Resources, Pollution and Society, William W. Murdoch, Ed., Sinauer Ass., Stamford, Conn.

Posner, G., 1954: The Peru Current, <u>Scientific American</u>, Vol. 190; 66-71.

Schweigger, E. H., 1931: Observaciones Oceanograficas Sobre la Corriente de Humboldt, Bol. Comp. Admin Guano., 7, (1):3-39.

Smith, R. L., 1978: Poleward Propagating Perturbations in Currents and Sea Level Along the Peru Coast, J. Geophys. Res., 83: 6083-6092.

Sverdrup, H. U., 1930: Some Oceanographic Results of the Carnegie's Work in the Pacific, The Peruvian Current, <u>Trans. Amer. Geophys., Un.</u>, 10th-11th An. Meeting.

Tessan, U.De., 1844: <u>Cartes des Courants et des Temperatures de l'Eau a la Surface de la Mer, Observes a Bord de la Venus en 1837, 38 et 39, Paris.</u>

Wooster, W., and Gilmartin, M., 1961: The Peru-Chile Undercurrent, \underline{J} . Mar. Res., 19 (13):97-122.

Wyrtki, K., 1963: The Horizontal and Vertical Field of Motion in the Peru Current, <u>Bull. Scripps Instn. Oceanogr.</u>, 8 (4):313-346.

Wyrtki, K., 1975: El Nino: The Dynamic Response of the Equatorial Pacific Ocean to Atmospheric Forcing, J. Phys. Oceanogr., 5: 572-584.

Wyrtki, K., 1973: Teleconnections in the Equatorial Pacific Ocean, Science, 183: 66.

Yergen, W., 1970: The Data Base of the Naval Oceanographic Live Atlas, Informal Report #70-1 NAVOCEANO.

Yergen, W., 1967: A Rapid Access Tape Format for Oceanographic Station Data, Informal Report #67-X NAVOCEANO.

Zarate, A., 1555: Historia del Descubrimiento y Conquista del Peru, con las Cosas Naturales que Senaladamente Alli se Hallan y los Sucesos que ha Avido, Anvers. (Traducido en Kerr, Roberts, 1824-), A General History and Collection of Voyages and Travels, IV:348-350.

INITIAL DISTRIBUTION LIST

		No.	Copies
1.	Defense Technical Information Center Cameron Station Alexandria, VA 22314		2
2.	Library, Code 0142 Naval Postgraduate School Monterey, CA 93940		2
3.	Chairman (Code 68Mr) Department of Oceanography Naval Postgraduate School Monterey, CA 93940		1
4.	Chairman (Code 63Rd) Department of Meteorology Naval Postgraduate School Monterey, CA 93940		1
5.	TNT Hector Soldi Direccion de Hidrografia y Navegacion de la Marina Esquina de Roca y Gamarra Chucuito Callao, Peru		15
6.	Director Naval Oceanography Division Naval Observatory 34th and Massachusetts Avenue NW Washington, D.C. 20390		1
7.	Commander Naval Oceanography Command NSTL Station Bay St. Louis, MS 39522		1
8.	Commanding Officer Naval Oceanographic Office NSTL Station Bay St. Louis, MS 39522		1
9.	Commanding Officer Fleet Numerical Oceanography Center Monterey, CA 93940		1

		No.	Copies
10.	Commanding Officer Naval Ocean Research and Development Activity NSTL Station Bay St. Louis, MS 39522		1
11.	Commanding Officer Naval Environmental Prediction Research Facility Monterey, CA 93940		1
12.	Chairman, Oceanography Department U.S. Naval Academy Annapolis, MD 21402		1
13.	Chief of Naval Research 800 N. Quincy Street Arlington, VA 22217		1
14.	Office of Naval Research (Code 480) Naval Ocean Research and Development Activity NSTL Station Bay St. Louis, MS 39522		1
15.	Scientific Liaison Office Office of Naval Research Scripps Institution of Oceanography La Jolla, CA 92037		1
16.	Library Scripps Institution of Oceanography P.O. Box 2367 La Jolla, CA 92037		1
17.	Library Department of Oceanography University of Washington Seattle, WA 98105		1
18.	Library CICESE P.O. Box 4803 San Ysidro, CA 92073		1
19.	Library School of Oceanography Oregon State University Corvallis, OR 97331		1

		No. Copies
20.	Commander Oceanographic Systems Pacific Box 1390 Pearl Harbor, HI 96860	1
21.	Chief, Ocean Services Division National Oceanic and Atmospheric Administration 8060 Thirteenth Street Silver Springs, MD 20910	1
22.	IHO/FIG International Advisory Board International Hydrographic Bureau Avenue President J. F. Kennedy Monte-Carlo, Monaco	1
23.	Hydrographer of the Navy Ministry of Defense Hydrographic Department Taunton, Somerset TA129N England	1
24.	Hydrographer, Royal Australian Navy Hydrographic Office Box 1332 North Sydney 2060 New South Wales - Australia	1
25.	Dominion Hydrographer Canadian Hydrographic Service 615 Booth Street Ottawa, Ontario KiAOE6	1
26.	Direccion General De Oceanografia Departmento De Hidrograficia Ave COYOACAN 131 Mexico 12, D.F.	1
27.	Director Instituto Hidrografico Rua Das Trinas, 49 Lisboa Portugal	1
28.	Director De Hidrografia y Navegacion de la Marina Direccion de Hidrografia y Navegacion de la Marina Calle Saenz Pena -5ta Cuadra-La Punta Callo-5 Peru	1

		No.	Copies
29.	Doug McLain PEG, NMFS P.O. Box 831 Monterey, CA 93942		1
30.	David Husby PEG, NMFS P.O. Box 831 Monterey, CA 93942		1
31.	Professor Roland Garwood Naval Postgraduate School Department of Oceanography Monterey, CA 93940		1
32.	LCDR Donald D. Winter NOAA Ships RUDE & HECK Atlantic Marine Center 439 W. York Street Norfolk, VA 23510		1
33.	LT Virginia E. Newell NOAA Ship DAVIDSON 1801 Fairview Avenue, East Seattle, Washington 98102		1
34.	Mr. Richard Joy DMAHTC - HYHO4 6500 Brookes Lane Washington, D.C. 20315		1
35.	TN Antonio Ruiz Instituto Hidrografico de la Marina Cadiz, Spain		1
36.	LT Laurent Monsaigeon Etat-Major de la Marine (EMM/MAT/SSM) 2. Rue Royale 75200 Paris Naval France		1
37.	LT Vitor Silva Ave. Forcas Armadas, Lote: A, Y, L, R; 1-ESQ 1600, Lisbon, Portugal		1
38.	Gene and Mary T. Brown Code 7300 Naval Oceanographic Office NSTL Station Bay St. Louis, MS 39522		1

No. Copies

1

39. Patricia Eaton
DMAHTC - HYHO2
6500 Brookes Lane
Washington, D.C. 20315

